CA20NUF.359P41 ENG

a31188003958077b

WAR MEMORIAL DOORWAY

This book is the property of
MacNabb Memorial Library
Ontario Veterinary College

CA2ΦNUF3

59P41

SF 756.T

P697

1959

THE UNIVERSITY OF TORONTO LIBRARY

MANUSCRIPT THESIS

AUTHORITY TO DISTRIBUTE

AUTHOR: This thesis may be loaned or individual microfilm copies made available:

(signature of author in *one* of
the three spaces below)

(a) without restriction

OR (b) with the restriction that,
for a period of five years (until)
the written approval of the
head of the graduate de-
partment is required.

OR (c) with the restriction that,
for a period of five years (until)
the written approval of the
author is required.

BORROWERS: The borrower undertakes to give proper credit for any use made of the thesis, and to obtain the consent of the author if it is proposed to make extensive quotations, or to reproduce the thesis in whole or in part.

[illegible]

METABOLISM OF PHENOTHIAZINE IN CHICKENS

by

Nicolas Platonow

A Thesis

Presented to the Faculty of the Graduate School
of the University of Toronto in Partial Fulfilment
of the Requirements for the Degree of Master of
Veterinary Science.

1959.

BIOGRAPHICAL SKETCH

The author was born November 30, 1928 in Lipscha, Czechoslovakia, where he attended his primary school and the lyceum. In May, 1957 he graduated from the University of Montreal with the Degree of D.M.V. In September, 1957 he enrolled in the School of Graduate Studies at the University of Toronto and for two years worked under the direction of Professor W. T. Oliver of the Division of Pharmacology at the Ontario Veterinary College, Guelph, Canada, towards the degree of Master in Veterinary Science.

ACKNOWLEDGMENTS

The author wishes to express his appreciation and thanks to Dr. W. T. Oliver for his constant guidance and helpful advise^c throughout the course of this study, and for constructive criticism during the preparation of this thesis.

Thanks are due to Mr. H. Funnel for his technical collaboration, to Mr. E. Dillistone for production of the graphs, to Dr. H. Neely for his aid in the photography, and to Mrs. J. Henry for her tireless typing and retyping of this manuscript.

TABLE OF CONTENTS

	Page
INTRODUCTION AND REVIEW OF LITERATURE	1
EXPERIMENTAL PROCEDURE	
Purification of Phenothiazine and Production of its Oxidation Derivatives.	3
Chromatography and Spectrophotometry of Phenothiazine and its Derivatives.	8
Chromatography and Spectrophotometry of Phenothiazine and its Derivatives after Extraction from the Body.	13
Isolation of the Metabolites from Bile.	14
Isolation and Identification of Phenothiazine and its Metabolites in the Caecal Contents.	15
Isolation and Identification of Phenothiazine and its Metabolites in Excreta.	16
RESULTS	19
DISCUSSION AND CONCLUSIONS	26
SUMMARY	32
BIBLIOGRAPHY	33
TABLES	38
FIGURES	49

INTRODUCTION AND REVIEW OF LITERATURE

Thiodiphenylamine (phenothiazine) was first prepared by Bernthsen (4) in 1883 as a base for colouring materials (5). Carro's (8) discovery in 1876 of methylene blue, one of phenothiazine's derivatives, gave Ehrlich a relatively successful antipaludic agent. This event opened the era of chemotherapy, and phenothiazine was forgotten.

During an investigation of the biological effects of sulfur compounds on insects, Campbell et al (7) found that phenothiazine had insecticidal properties. In 1938 Knipling (35) used phenothiazine as a treatment for ectoparasites of cattle, and in the same year Harwood et al (32) reported on its anthelmintic properties. In the next two years its anthelmintic properties in man and domestic animals were investigated.

The first reports of the effects of the administration of phenothiazine to poultry appeared in 1940 (43 and 44) and were concerned with its vermifugal activity. Since that time one group of workers published a series of articles on the effects of phenothiazine on the uptake of I^{131} by the thyroid gland of various animal species including poultry (63) but the remainder of the reported in vivo studies on phenothiazine in poultry have been limited to its use as an anthelmintic against Heterakis gallinae and Ascaridia galli (1,24,25,26,30,31, 41,42,47,50,54 and 61) Capillaria spp. (22,37,56 and 60) and its use in the prevention and treatment of Syngamus trachea infections (9). Avian coccidiosis was also reported to have been cured, and outbreaks

of the disease controlled by the prophylactic and curative use of the drug (18,57 and 62); in powder form it was found to be useful in the control of louse infestations of chickens (52). Determinations of the drug's toxicity under field conditions were also undertaken (44 and 45).

In vitro studies were conducted to investigate the metabolism and pharmacological actions of phenothiazine and its derivatives in Heterakis gallinae (48) and the fate in nematodes of phenothiazine labelled with S^{35} have also been reported (21 and 38).

Since phenothiazine possesses one of the widest therapeutic spectra of the anthelmintics, it is still one of the most widely used vermifuges, consequently its pharmacodynamics is of considerable importance. After a recent and comprehensive survey of the literature, Harwood was led to report (29):

"The fate of phenothiazine has been determined in general terms, but many details are still in doubt or disputed because satisfactory methods of analysis are not standardized, particularly methods which will distinguish between phenothiazine and each of the several compounds that may be derived from it by oxidation."

The experiments reported herein were conducted in order to study the problem of the metabolism in chickens of purified forms of this drug.

EXPERIMENTAL PROCEDURE

Purification of Phenothiazine and Production of its Oxidation Derivatives.

The following procedures were undertaken to obtain purified phenothiazine and its oxidation derivatives for administration to the experimental chickens, comparison purposes during paper chromatography and spectrophotometric analyses, and for the investigation of colorimetric methods for quantitative determinations of the drug and its metabolites.

(a) Phenothiazine. (Thiodiphenylamine, Imido-diphenyl-sulfide)

Phenothiazine N.F. (green)* was purified with charcoal** by recrystallization from 95 per cent ethanol, as follows:

Phenothiazine N.F. (green) was treated with Norit-A at a concentration of 5 per cent of the weight of phenothiazine. The boiling saturated solution of phenothiazine in 95 per cent ethanol was filtered quickly through Whatman No. 1 into a heated glass funnel. The filtrate was kept overnight in a freezer at a temperature of -10 to -12°C . The following morning the crystals were collected by filtration. The procedure was repeated three times. After distillation the used ethanol was utilized again for the first two steps of the purification.

The final product yielded pale yellow crystals, with a melting point of: 183.5 to 184.5°C .*** Yields of 60-65 per cent were obtained.

* Phenothiazine N.F. (green): Fisher Scientific Company.

** Charcoal: Norit-A (Decolorizing Carbon) Neutral,
Fisher Scientific Company

*** M.P. $182.5 - 183.5^{\circ}\text{C}$ reported by Houston et al (34)

(b) Diphenylamine - ortho - Sulfoxide. (Phenothiazine-5-oxide)

Phenothiazine-5-oxide was prepared by the method of Pummerer and Gassner (53). The yield was 70 per cent (based on phenothiazine) and the product had a melting point range of 251 to 252°C*. The crystals were very slightly yellow and had polygonal-cylindrical forms, with an average diameter of 30 μ and average length of 300-400 μ .

(c) Phenothiazone-3. (3-Hydroxyphenothiazine; Phenothiazone)

Phenothiazone-3 was prepared by the sulfur** fusion process of p.-hydroxydiphenylamine (33). The dried and ground solids so obtained were washed with 0.5 per cent potassium hydroxide, followed by cold water. They were then purified as follows:

Crude phenothiazone-3 was dissolved in chloroform (purified by fractionated distillation) and filtered through a column of aluminium oxide powder***. The major part of the collected eluate was evaporated in a water-bath, and the evaporation was completed in the vacuum dryer, using a mechanical pump. Further purification was achieved by crystallization from 20 per cent diluted ethanol. The product was brick-red, with a melting point of 162-163°C****; its crystals had tetragonal-cylindrical forms, with an average diameter of 5 μ and an average length of 10-20 μ .

The p.-hydroxydiphenylamine (p.-anilinophenol) used was obtained from a commercial source*****; it was purified by two successive

* M.P. 250°C reported by Harpur *et al* (28) and by Barnett *et al* (2)

** Sublimed sulfur (U.S.P.): Fisher Scientific Company

*** Aluminium Oxide Powder: Baker Analyzed Reagent

**** M.P. 161°C reported by Houston *et al* (34)

***** p.-Anilinophenol (Technical): Eastman Kodak Company

distillations with superheated steam, and the final product consisted of pearly flakes with a melting point of 69-70°C*.

Ultraviolet absorption of phenothiazine, phenothiazine-5-oxide and phenothiazone-3 in absolute methanol at the concentration 5×10^{-5} M have been measured, with Beckman DU-Spectrophotometer, for control and comparison purposes (Figure No. 6). Phenothiazine: maxima at 255 and 320 m μ , inflections at 235-240 and 280-285 m μ . Phenothiazine-5-oxide: maxima at 230, 270, 305 and 340 m μ , inflections at 250, 285 and 320 m μ . Phenothiazone-3: maxima at 237-238, 273 and 370 m μ , inflections at 250 and 320 m μ .

(d) Thionol (7-Hydroxyphenothiazone-3)

(i) The first attempt at thionol production was made according to the method of De Eds et al (13). A dark-blue amorphous mass was obtained, which did not melt at 360°C. This was dissolved in boiling glacial acetic acid and gave an intense red color. By paper chromatography of this solution, we obtained two distinct spots, corresponding to the R_f value of phenothiazine and phenothiazone.

(ii) By the original Granick and Michaelis (23) method the yield was insignificant.

(iii) More interesting results were obtained by the sulfur fusion of p.,p'-dihydroxydiphenylamine process of Houston et al (34). The original process gave a dark-blue amorphous mass, which did not melt

* M.P. 70°C reported by Calm (6)

at 315°C and whose final yield was low. The paper chromatogram of this product dissolved in glacial acetic acid, gave several separate spots, which indicated impurities.

By carrying out the following modification of the above process, we obtained a purple amorphous mass, which did not melt at 360°C and which by chromatographic analysis was almost pure.

After fusion of equivalent amounts of sulfur and p.,p'-dihydroxydiphenylamine, with iodine as catalyzer, at 195-200°C for 60 minutes, the residue was removed and washed with sodium hydroxide 0.5 per cent, followed by cold water. The dried product was dissolved in ethanol 95 per cent, and this solution was filtered twice through a column of aluminium oxide powder (1 x 10 in. high). Evaporation of ethanol eluate was commenced on a hot plate, continued in a water-bath, and completed in a vacuum dryer at 37°C in darkness. The residue was then dissolved in acetone and passed through an alumina column. The thionol was adsorbed by aluminium oxide, and the phenothiazone was passed through the column with acetone. The thionol which was absorbed by the alumina column was removed from it by the addition of ethanol (95 per cent), and the ethanol eluate was evaporated in the manner described above.

The molar extinction coefficient of thionol obtained in this way was 61,500* in 0.01 N sodium hydroxide at its maximum absorption at 590 mμ, and 32,400 in 8N sulfuric acid at its maximum absorption at 500 mμ (Figure No. 7).

*63,000 reported by Cranick and Michaelis (23).

p.,p'-Dihydroxydiphenylamine was prepared according to Knoevenagel (36) and melted at 167-168°C (Knoevenagel reported 169°C).

(iv) Thionol in crystalline state was obtained by the following modification of the Granick and Michaelis (23) method:

Digestion time was reduced to 6 hours at a temperature of 160-165°C, using 80 per cent (by volume) sulfuric acid (325 ml of 80 per cent sulfuric acid for 22 gms of phenothiazine N.F.(green).

Yield was 7.5 per cent* of thionol (based on phenothiazine), in the form of an agglomeration of tetragonal brown-reddish needles with constant face area (15 μ x 15 μ). The crystals did not melt at 360°C.**

* 5 per cent reported by Granick and Michaelis (23).

** idem reported by Houston et al (34).

Chromatography and Spectrophotometry of Phenothiazine and its Derivatives.

A. Paper Chromatography

a) Apparatus.

Ascending paper chromatography was used for the separation of phenothiazine and its oxidation derivatives from various body extracts and of phenothiazine from its derivatives. By the technique herein described good separation of the fractions was obtained without oxidative changes of the metabolites during the separatory procedure.

The apparatus consisted of a chromatographic chamber (Figure No. 1) and its covering apparatus (~~Figure No. 2~~), a vacuum pump*, and a nitrogen bomb**.

The chromatographic chamber was an 18 by 13 by $\frac{1}{4}$ inches Pyrex cylinder fitted above and below with $\frac{3}{4}$ inch thick stainless steel plates, 15 inches in diameter. Taking 50,000 p.s.i. as the ultimate compression stress and using a safety factor of 5, from the formula

$$(51) \quad T = \frac{(F \times P) + 1386 D}{86670},$$

the Pyrex cylinder used in these experiments with a wall thickness of 0.25 of an inch will withstand maximum safe pressure of 91 p.s.i. The upper plate was drilled and tapped at its center to allow the insertion of a teflon stopcock. This stopcock was used for the evacuation of air from the vessel and the addition of nitrogen gas. To improve contact between the plates and the rims of the cylinder, polyethylene sheets were fixed on the inside

* Duo-Seal Vacuum Pump: W. M. Welsh Manufacturing Co., Chicago, U.S.A.

** Dry Nitrogen: The Lindae Air Products Co., Canada

the plates and the seal was made airtight by the use of vacuum grease. The polyethylene sheets also served to protect the steel plates from possible corrosion by the solvent. A polyethylene solvent trough, 12 inches in diameter and 1 inch high, was placed at the bottom of the chamber. (See: Figure No. 1)

A filter paper rack (Figure No. 1) for the rigid support of three paper sheets, 10 by 16 inches was made from two Pyrex glass frames, with outside dimensions 10 by 16 inches, and inside dimensions 8 by 14 inches. Four holes were drilled through the corners of the frames, and the frames were connected by four glass rods which passed through the holes and fastened with polyethylene washers. Four glass tubes, $1\frac{1}{2}$ inches long, were threaded on each rod to serve as spacers. The tubes prevented the paper sheets from sliding along the rods and also ensured that the sheets were maintained equidistant from one another. The stopcock on the upper plate was connected by means of rubber tubes to a vacuum pump and nitrogen bomb.

To prevent photochemical changes of phenothiazine and its metabolites during chromatography, the entire chromatographic chamber was covered with a bottomless box, made from one-half inch thick plywood, 21 inches long, 21 inches wide and 24 inches high (Figure No. 2). A hole, one-half inch in diameter, was made in the center of the top for passage of the rubber tube from the vacuum pump to the teflon stopcock. A window 12 by 12 inches was made at the front of the box for convenient observation of the progress of the solvent front.

b) Method.

On a sheet of filter paper* 10 by 16 inches, four pencil marks were made 2 inches apart and $1\frac{1}{2}$ inches from the lower edge of the sheet. Samples of the drug and its metabolites were applied to the marks with micropipettes.

To increase the definition of separation of each fraction, the chamber was saturated with water vapour prior to chromatography. Two glass mortars with distilled water were placed inside the solvent trough, and the inner wall of the glass cylinder was covered with filter paper moistened in distilled water.

A mixture of acetic acid, benzene and water in the proportions 200: 400: 600 mls. was used as the developing solvent.

When the chamber was saturated with water and solvent vapour (this equilibration process took at least 16 hours), the rack holding the filter papers with the samples was placed in a vertical position in the solvent trough and the cabinet was made airtight. A rubber tube joining the vacuum pump and the nitrogen bomb was then connected to the stopcock situated on the upper steel plate. Air from the chamber was drawn out by the vacuum pump and nitrogen was then introduced until the pressure inside the cylinder was identical with that of the atmosphere. The same procedure was repeated three times in order to ensure the complete evacuation of air from the cylinder and the establishment of anaerobic conditions. Sixteen hours were required to obtain a good chromatographic separation.

* Whatman No. 1: Fisher Scientific Co., Toronto

The chromatograms were dried in an oven at 50° for 30 minutes and then sprayed with a one percent alcoholic solution of iodine to locate the limits of the fractions, though the positions of thionol and phenothiazine were usually visible without spraying. The spot of phenothiazine-5-oxide was more easily detected when sprayed with hydrochloric acid 8N.

A typical chromatogram of phenothiazine and its oxidation derivatives is given in Figure No. 3.

B. Spectrophotometry.

1) Qualitative determination of phenothiazine, phenothiazine and thionol was established by the method of Oliver and Funnell (49).

2) The colorimetric determination of phenothiazine-5-oxide.

According to Barnett and Smiles (2) when the sulfoxide is submitted to the action of aqueous hydrochloric acid it is at once converted into phenazothionium chloride which is a stable salt. Diphenylamine-ortho-sulfoxide is colourless in acetic acid in the visible absorption range, but with the addition of hydrochloric acid the solution becomes red-brown and gives the highest absorption peak at 520mμ (Figure No. 4). Although the change is favoured by increasing the concentration of the acid, the concentration of 8N is found to be most convenient to handle.

After chromatographic separation the section of paper containing the phenothiazine-5-oxide was removed and put into a 50 ml. Erlenmeyer flask. The product was eluted from the paper by the addition of 5 ml. of glacial acetic acid, the flask being stoppered and agitated for 30

minutes. The solution was filtered and a three ml. aliquot was transferred to another 50 ml. flask and three ml. of an 8N solution of hydrochloric acid containing one drop of a one percent solution of ferric chloride in 8N hydrochloric acid were added. The flask was stoppered and agitated for two hours.

The solution was read photometrically against a blank prepared by the same method as the unknown.

Determination of the calibration curve of phenothiazine-5-oxide:

Exactly 50 mgms. of phenothiazine-5-oxide were dissolved in 100 ml. of glacial acetic acid. Ten ml. of this solution were pipetted into a 200 ml. volumetric flask and diluted to 200 ml. with glacial acetic acid.

Two, five, 10, 15, 20 and 25 mls. of the diluted solution were pipetted into separate 50 ml. Erlenmeyer flasks. To each solution were added respectively, 23, 20, 15, 10, 5 and nil mls. of glacial acetic acid. The vessels were stoppered and agitated on a mechanical mixer for 15 minutes. Fifteen mls. of each solution were transferred to a second 50 ml. Erlenmeyer flask and 15 ml. of 8N hydrochloric acid and 5 drops of a one percent solution of FeCl_3 in 8N hydrochloric acid were added to each sample. The solutions were stoppered and agitated mechanically for two hours.

They were then read in the Klett-Summerson photometer, using a No. 52 filter, against a blank prepared by adding 15 ml. glacial acetic acid, 15 ml. HCl (8N) and 5 drops FeCl_3 (1% in 8N HCl). The reaction followed Beer's Law for the concentrations tested (Figure No. 5).

Chromatography and Spectrophotometry of Phenothiazine and its Derivatives after Extraction from the Body.

DeKalb cockerells*, 8 to 10 weeks old were used in this study. These birds had been on commercial mash prior to experiment. Chickens to be used for drug metabolic studies were fasted overnight and dosed with one gram of purified phenothiazine by means of gelatin capsules administered orally and manipulated directly into the crop.

Each bird was killed at a designated time after drug administration. The time intervals selected were one-half, one, two, three, four, five, six, 16, 24, and 48 hours post-treatment.

Immediately after the bird was killed, the digestive tract, liver, kidneys and gallbladder were removed from the abdominal cavity. The digestive tract was divided arbitrarily into: crop, proventriculus, gizzard, small intestine (from gizzard to the junction of the caeca with the large intestine), caeca, large intestine and cloaca. The contents from the various parts of alimentary canal as well as excreta, homogenized liver, kidneys and gallbladder were transferred to Erlenmeyer flasks. The drug and its derivatives were extracted by an acetone-methanol-formaldehyde mixture (1:1:.05) which was added to the flasks at the rate of 10 ml. of solution for each gram of material to be extracted. The flasks were stoppered and mechanically shaken for 30 minutes and the extracts were filtered and kept in stoppered tubes in a dark place until analyzed.

* Harvey's Poultry Farm
R. R. 2, Guelph, Ontario

Fifty microliters of extract from the content of the crop, gizzard, proventriculus, small intestine, caeca, large intestine, and cloaca and from excreta, and 0.1 ml. of extract from liver and kidneys were applied to filter paper and chromatographic separation followed by quantitative determination of drug and its metabolites were performed as previously described. The determination of phenothiazine in serum was made by the methods of Harpur et al (27) and Collier et al (12); of leucophenothiazine in serum and plasma by the method of Harpur et al (27), and the total water soluble conjugates in plasma, kidneys and liver by a modified (20) method of Collier et al (12).

Isolation of the Metabolites from Bile.

During chromatography of bile extracts it was found that certain bile constituents interfered with the chromatographic separation of phenothiazine and thionol, producing diffuse, poorly defined patterns. To reduce this interference 0.5 ml. of bile extract was streaked $1\frac{1}{2}$ inches from the lower edge of the chromatographic paper. After chromatographic development the spots of drug metabolites were cut out and eluted with 25 ml. of absolute methanol. The eluate was evaporated to five ml. in an atmosphere of nitrogen. One ml. of the evaporated eluate was applied to a chromatographic sheet. Good separation of the metabolites was effected by the method described above.

Isolation and Identification of Phenothiazine and its Metabolites in Caecal Contents.

Caecal contents from a bird dosed with one gram of phenothiazine and killed 24 hours after drug administration were extracted with absolute methanol. The extract was filtered and transferred to a separatory funnel, twice the volume of reagent grade chloroform was added and the contents shaken vigorously. An excess of water was added to effect the separation of the chloroform solution. The chloroform phase was drawn slowly through an alumina column, previously moistened with chloroform.

An intense red band moved down the column and into the effluent and a blue-purple band was adsorbed on the top of the column. The column was washed with chloroform, followed by acetone to remove traces of unabsorbed substances. The chloroform eluate was evaporated to dryness, yielding a dark reddish-brown residue.

The residue was treated with a boiling water and, after cooling, was filtered. One-half of this water extract, hereafter referred to as Fraction No. 1, was treated with reagent grade chloroform and benzene, to permit transfer of this coloured material to those particular solvents. The other half was divided into three portions; d-tartaric acid was added to one portion, acetic acid to the second and sodium hydroxide to the third, in order to obtain half saturated, 50 per cent, and 0.01 N solutions respectively.

The residue remaining after water treatment was washed several times with boiling water and recrystallized by the method previously described (See: Purification of phenothiazine). These crystals, here-

after referred to as Fraction No. 2, were dissolved in absolute methanol and the solution was read with spectrophotometer in the ultraviolet range.

The fraction adsorbed to the upper part of the alumina column was eluted from the column by the addition of 95 per cent ethanol and the eluate evaporated. The residue, which will be referred to as the Fraction No. 3, was divided into four portions. Each of those four portions were then dissolved with sodium hydroxide 0.01 N, sulfuric acid 8N, 95 per cent ethanol, and water respectively.

Each of above solutions was then scanned on the Beckman model DK spectrophotometer through the visual or ultraviolet spectrum ranges.

Isolation and Identification of Phenothiazine and its Metabolites in Excreta.

Three birds were fasted overnight and each was dosed with one gram of purified phenothiazine. The chickens were kept in metallic cages and the excreta was collected during the 24 hours period following drug administration.

The drug and its derivatives were extracted from the excreta with methanol, and the extract was evaporated on waterbath to one-fifth of its original volume. The concentrated methanol extract was transferred into a separatory funnel, twice its volume of chloroform was added and the two solutions were mixed by vigorous shaking. The addition of the chloroform precipitated the greater part of the methanol soluble excretory fractions other than phenothiazine and its metabolites which then were removed by filtration. The filtered mixture was transferred

into a large beaker and one-half its volume of distilled water was added. The two organic solvents were removed from the mixture by evaporation on a boiling waterbath. The removal of the organic solvents precipitated crude phenothiazine which was recovered by filtration. The coloured components in the aqueous phase then were extracted three times with chloroform.

The combined chloroform extract was drawn slowly through an alumina column, previously washed with chloroform. A blue-purple band appeared on the top of the column from which a reddish band separated and passed through the column with the eluate. The column was washed with chloroform and allowed to stand for a few hours at room temperature.

The chloroform eluate when evaporated on waterbath to dryness yielded a reddish-brown solid. The residue was recrystallized from 30 ml. of water and reddish crystals were recovered. The crude phenothiazine was purified by the method previously described (See: Purification of phenothiazine).

The blue-purple band which had adsorbed on the top of the column, was removed by the addition of 95 per cent ethanol, leaving green and grey bands at the top of the column. By rechromatography of this ethanol eluate on a fresh column washed with ethyl alcohol, only a trace of green and grey bands appear on the top of the column. After filtration the ethanol eluate was evaporated to dryness.

The melting points, ultraviolet, infra-red and visible transmittances of ethanol-eluate residue (Fraction B), crystals from water solution (Fraction A), and those obtained by three times recrystallization in 95 per cent ethanol (Fraction C) were measured. The determination of glucuronic acid in Fractions A and B was made by a modified (3) method of Mozolowski (46).

RESULTS

The distribution of phenothiazine and its oxidative metabolites, phenothiazone and thionol, in the contents of the digestive tract and excreta, and in the liver, bile, kidneys and plasma, was established as previously specified, and the data are presented in Tables 1, 2, 3 and 4.

Crop. Phenothiazine reached its maximum concentration in the crop, 835 mgms, 30 minutes after administration. The concentration of the drug then decreased constantly with time, so that 24 hours after treatment only 1.3 mgms were detectable. At 48 hours after the dosing no detectable amount was present in this part of the digestive tract.

Proventriculus. There was no evidence of extensive accumulation of the drug in the proventriculus. The phenothiazine level in this portion of digestive tract was low throughout the experiment. It reached a concentration of 5.7 mgms at 30 minutes, increased to 23 mgms at two hours post-treatment, decreased to 1.6 mgms at three hours, and remained relatively unchanged from the third to the 24th hour after treatment. No drug could be detected in the proventriculus 48 hours after dosing, thus its disappearance from this organ corresponded with its removal from the crop.

Gizzard. Phenothiazine appeared in the gizzard shortly after its administration to the birds. Its concentration in this organ was 15.2 mgms within the first 30 minutes and the drug reached its peak, 470 mgms, in the gizzard at four hours post-treatment. After four hours

there was a consistent decrease in the drug concentration with time. Only 6 mgms were present at 48 hours.

Neither the crop, proventriculus nor the gizzard contained detectable quantities of the oxidative derivatives, even though large quantities of the parent compound were present.

Small intestine. Three mgms of phenothiazine were detected in the small intestine one hour after administration. The drug concentration in this portion of the intestines increased slowly, reaching its maximum concentration of 330 mgms at six hours. The concentration remained high until 16 hours after dosing, and decreased slowly during the following 32 hours.

Phenothiazone was first recovered from the small intestine at six hours, and thionol at 16 hours post-treatment. The amount of phenothiazone was greater than thionol at 16 hours, lower at 24 hours, but their concentrations were essentially equal at 48 hours. Minimum levels of each of the metabolites in the small intestine, 0.36 mgm of phenothiazone and 0.34 mgm of thionol, occurred at 48 hours.

Caeca. Phenothiazine and its metabolites which were isolated by means of an alumina column (See: Experimental Procedure) from the caecal contents were analyzed on the Beckman Model-DK spectrophotometer. Three fractions were recovered. In the visual spectrum, Fraction No. 1 gave the same maxima and inflections as synthetic phenothiazone in five solvents. The results are presented in Table No. 5 and in Figures Nos. 8,9,10,11,12 and 13.

In the ultraviolet spectrum with absolute methanol as solvent Fraction No. 2 (Figure No. 14) gave minima at 253 and 316 mμ and the maxima at 281-282 mμ. A similar curve was obtained using purified phenothiazine.

In the visible spectrum, using four different solvents, Fraction No. 3 gave minima which were identical with synthetic thionol when calculated by the Lewis rule (40). However, in each case the maximum was shifted to the left. The results are presented in Tables No. 6 and Figures Nos. 15, 16, 17 and 18.

Phenothiazine was first detected in the caeca at the fourth hour, and its concentration in this part of the tract coincided with the progression of the drug through the small intestine. It increased constantly during the succeeding hours reaching its peak, 15 mgms, at 24 hours. Its removal from this site was slow, 40 per cent (6.7 mgm) of the maximum concentration attained still remained at 48 hours.

Phenothiazone was first detected in the caeca at four hours and thionol at five hours post-treatment. Maximum concentrations of phenothiazone and thionol, 3.0 and 0.9 mgms, respectively, occurred at 16 hours. After the 16th hour the levels decreased, but were relatively high 48 hours after the administration of the parent compound. Large intestine. In the large intestine phenothiazine was detected at four hours. It reached its maximum concentration, 0.8 mgm, by the 16th hour and its level then decreased slowly during the following 32 hours. 0.1 mgm were present 48 hours after administration.

Phenothiazone and thionol appeared in large intestine at five hours post-treatment. The amount of phenothiazone was consistently greater than that of thionol throughout the duration of the experiment. Maximum level of phenothiazone was at 48th hour, and those of thionol at 16th hour. Both metabolites were present in large intestine in detectable amounts 48 hours after treatment.

Cloaca and Excreta. Phenothiazine and its oxidative metabolites isolated from excreta in crystalline forms (See: Experimental Procedure) were analyzed on the Beckman Model-DK spectrophotometer in the ultraviolet and visible transmittance ranges and on the Beckman Model-IR 4 spectrophotometer in the infra-red range. Fraction A gave the same maxima and minima as synthetic phenothiazone. The results are presented in Table No. 7 and Figures Nos. 19 and 20. In the ultraviolet and visible ranges using three solvents Fraction B gave the same minima as synthetic thionol. The results are presented in Table No. 8 and Figures Nos. 21, 22, 23 and 24. By the Mozolowski test (46) Fraction A was negative but Fraction B contained 1.02 per cent of glucuronic acid. In the ultraviolet range in absolute methanol Fraction C gave a pattern identical with pure phenothiazine. The results are presented in Table No. 9 and Figure No. 25. The infra-red spectra of the Fraction A (Figure No. 26) and Fraction B (Figure No. 28) in contrast to that of synthetic phenothiazone (Figure No. 27) and thionol (Figure No. 29) were apparently identical.

Phenothiazine was detected first in cloaca and excreta at four hours post-treatment, and at five hours over 10 per cent of the

original dose of the drug administered to the birds was present. The excretory rates increased constantly and at 48 hours the greater portion of the drug had been eliminated. The phenothiazine isolated from excreta in crystalline form (See: Experimental Procedure) gave following melting point: 182.5°C . When this purified excretory product was mixed with a purified synthetic phenothiazine, no depression in melting point was observed.

The first appearance of thionol in cloaca and excreta was within the first hour post-treatment. At three hours the second metabolite, phenothiazone, appeared in cloaca and at four hours in the excreta. After four hours the amounts of phenothiazone and thionol in cloaca and excreta increased constantly with time and reached its maxima at 48 hours after administration.

Bile. After rechromatography of the metabolites (See: Experimental Procedure) the upper and lower spots whose R_f values corresponded to phenothiazone and thionol respectively, were eluted from paper with absolute methanol. The solutions were read on Beckman Model-DU spectrophotometer in the visual range.

The absorption maxima of the upper spot eluate occurred at 370 and 505 $\text{m}\mu$ with an inflection at 420 $\text{m}\mu$. The same pattern was observed with synthetic phenothiazone. The eluate from the lower spot gave absorption maxima at 425 and 590 $\text{m}\mu$, with an inflection at 450 $\text{m}\mu$. Similar results were recorded with synthetic thionol.

Phenothiazine was first recovered in the bile four hours after dosing. The concentration of the drug increased from 0.15 mgm to a

maximum of 0.4 mgm six hours after administration. Only a small amount (0.06 mgm) remained in the gallbladder at the 48th hour.

The oxidation derivatives, phenothiazone and thionol, appeared simultaneously at the fifth hour. The concentrations of phenothiazone were twice as much as those of thionol at the fifth, 16th and 24th hours, practically equal at the sixth hour and lower at the 48th hour.

Liver. Although the drug was detectable in this organ soon after treatment, (Figure No. 30), its concentration remained low throughout the experiment. The level from the first hour increased slightly, reaching its peak level at the fifth hour (1 mgm), after which it decreased slowly until the end of the experiment. At the 48th hour, the level of the drug was half of its maximum concentration (0.6 mgm) in this organ.

Water-soluble metabolites were observed within one hour of administration of the parent compound. The concentration increased from 0.05 mgm to a peak of 0.3 mgm at four hours. Only 0.08 mgm were present after 16 hours. The level of the fractions in the liver after the fourth hour decreased in a parallel fashion with the appearance in the bile of phenothiazone and thionol. At the 24th hour the amount of metabolites increased to 0.12 mgm, increasing to 0.15 mgm by the 48th hour.

Kidneys. No phenothiazine could be detected in the kidneys during any stages of the experiment. The water-soluble oxygenated derivatives appeared in the kidneys within the first hour after drug administration coinciding with the appearance of thionol in the cloaca and excreta. The level at the end of one hour, which was 0.1 mgm, fluctuated throughout

the remainder of the experiment. Thus the second and fifth hour levels were higher than the first, third, fourth and sixth hour levels respectively. The 16th hour concentration was lower than the sixth hour but higher than the 24th hour, which was lower than the 48th hour concentration.

Plasma. No diphenylamine ortho-sulfoxide was detected by the method of Harpur et al (27) in the serum or plasma of the dosed bird at any time during the experiment.

The presence of phenothiazine and leucophenothiazone in the serum and in the plasma was established by the first hour. The amounts of phenothiazine and its total oxygenated derivatives at that time were 0.3 and 0.6 mgm per 100 ml. of plasma respectively. The concentration of phenothiazine increased during the following hours, reaching its level, 3.5 mgms, at the 16th hour. By the 48th hour, the amount of drug present in the circulation was slightly lower than at the end of the first hour. The oxidation derivatives in plasma were higher in concentration but parallel to that of the parent compound throughout the experiment.

DISCUSSION AND CONCLUSIONS

The study of the metabolism of purified phenothiazine in the chicken was investigated by means of paper chromatography, spectrophotometry and by direct colorimetric methods.

It was evident one hour after administration that in the bird, as in the rat (15 and 16), rabbit (12 and 15), swine (12) and man (15,16 and 17), the absorption of phenothiazine could proceed independently of any detectable oxidative change in the molecule. Attempts to devise a surgical technique for the collection of blood from the portal-hepatic venous system were unsuccessful, thus it was not possible to establish directly whether the compound was absorbable only as phenothiazine. However, since

- (1) only phenothiazine was present in the small intestine, when only phenothiazine and leuco-phenothiazone could be detected in serum and plasma, and
- (2) only phenothiazine was present in the small intestine when phenothiazine and its derivatives were also detected in the liver, it was concluded that all of the compound absorbed during the first four hours following administration was absorbed as phenothiazine.

From the fifth to the 48th hour the concentration of the oxidative derivatives in blood and bile varied directly with their concentration in the lower intestinal tract. Throughout this time, since phenothiazine derivatives were detected only adjacent and

posterior to the bile duct, it appeared that any derivatives present in the intestinal tract were derived by way of the bile and blood from that portion of the phenothiazine which had been absorbed and had undergone metabolic oxidation.

During this same period the plasma concentration of phenothiazine varied directly with the concentration of the drug in the small intestine. Conversely, the concentration of oxidative derivatives in the plasma varied inversely with their concentration in the small intestine. Since no absorptive mechanism has been described in which the rate of intestinal absorption of a drug increases with a decreasing concentration gradient, the previously described observations were supported and it was concluded that in the bird, unlike calves and pigs, only phenothiazine was absorbed.

As stated previously, because of the technical difficulties of collecting blood draining the alimentary tract, particularly the small intestine, of poultry, direct analyses were impossible. However, the presence of phenothiazine in the serum and plasma was detected only after the drug had reached the small intestine. Also, the concentration of phenothiazine in the plasma varied in the same direction as its concentration in the small intestine. This simultaneous detection of the drug in the small intestine and in the circulation and the comparable nature of the drug's concentration in the blood and intestinal tract gave good indirect evidence that the site of absorption is, at least chiefly, in the small intestine.

The metabolic fate of phenothiazine has been reported for various animal species (3,10,11,12,15,16,17,19,27,39 and 59) and man (15,16 and 17). The results indicated that, depending upon the species, the parent compound was converted, in part, to phenothiazone, thionol, their leuco-forms and/or diphenyl amine ortho-sulfoxide. In the chicken it was found that phenothiazine was oxidized to leucophenothiazone and leuco-thionol. There was no evidence, however, that this species can convert phenothiazine to diphenyl amine o-sulfoxide as reported for calves (11) and sheep (12).

The site of oxidation of phenothiazine has not been reported but since the oxidation products, phenothiazone and thionol were detected in the bile before either was present in the small intestine it gave good presumptive evidence that the liver was the initial site of oxidation. But the liver was not the sole site of oxidation. Phenothiazone was recovered from the caeca one hour before it was present in any other part of the alimentary tract indicating that phenothiazine was also being oxidized in these organs, an important area of anthelmintic activity of this drug in the bird. The result was interesting because it has been reported that phenothiazine also is oxidized in the caecum of horses (12) and rumen of cattle (10) and sheep (28), organs which are physiologically analogous to the caeca of the chickens (58). The observation also re-emphasized the possible relationship between bacterial action, oxidation and anthelmintic activity observed previously by De Eds and Thomas (17) and Oliver (48).

The major part of the phenothiazine which was administered orally to chickens in these experiments, was eliminated by way of the intestinal tract. Depending upon its metabolic fate, two excretory routes were available for the removal of the phenothiazine which was absorbed by the bird; the kidneys and the bile. The demonstration of phenothiazine in the bile indicated that the fraction of the absorbed phenothiazine which escaped oxidation was returned to the intestinal tract by way of the gallbladder. Since both thionol and phenothiazone were present in the bile before they were found in the small intestine, it was apparent that part of the oxidized fraction was also removed from the body by way of the bile. The remaining oxidized fraction was excreted through the kidneys. The failure to detect phenothiazine in either the kidneys or urethrae indicated that either the compound per se was not excreted by way of the urine or it was only excreted in insignificant amounts.

Phenothiazine was removed from the body in an exponential manner, that is, the rate of its removal at any instant was dependent upon the concentration of the drug in the body at that time. At one hour after administration, 95 per cent of the original dose remained in the body. There was little change in this concentration of phenothiazine during the ensuing first three to four hours, the time required for it to pass through the alimentary tract. Its removal thereafter was rapid. About 78 per cent was present at six hours; 30 per cent at 24 hours and approximately 12 per cent at 48 hours.

The derivatives, after reaching a maximum concentration between the sixth to 12th hour, also were excreted at a rate which depended upon the amount present. Approximately 40 per cent of the maximum level was excreted by 24 hours and only 1.04 mgm remained in the body at 48 hours.

Both phenothiazine and its derivatives were excreted most slowly from the caeca than any other site in the bird. Forty per cent of the maximum level of phenothiazine attained and 56 per cent of its derivatives remained at 48 hours after treatment. Thus in the caeca susceptible organisms are exposed to the action of phenothiazine or its metabolites for longer periods than parasitic forms in other sites, a factor which may explain in part, its pharmacological efficiency in this site.

Phenothiazine, when administered orally, to chickens was excreted from the body in four forms, unconjugated phenothiazine, unconjugated leucophenothiazone, unconjugated leucothionol and leucothionol conjugated with glucuronide. Both of the reduced forms were rapidly oxidized to phenothizone and thionol after passing from the cloaca.

The nature of the excretory products of this drug in other species are presented in Table No. 10. It is seen that the excretory forms in the chicken resemble those of the rabbit (3,12,15,16 and 17). Although the forms of the excretory products of phenothiazine were similar in the chicken and rabbit, the ratio of conjugated to non-conjugated thionol in the chicken was in the order of 1:99 whereas in the rabbit it was found to be 99:1.

In the chicken the excreted fraction differed from most other species investigated chiefly in the nature of the conjugated fraction. In our case spectral analyses of extracts at various times and from various tissues and excreta failed to reveal the presence of the sulfate radical which was common to the conjugated excretory forms in the sheep, horse and cow.

From these results it appears that if the bio-activity of phenothiazine depends upon some metabolic change by the body, the pharmacologically significant fraction is either leucophenothiazone or leucothionol, results which supported previously reported in vitro investigations of the pharmacological mechanism of phenothiazine (48).

SUMMARY

Methods are given for the purification of phenothiazine N. F. and the preparation of its oxidation derivatives, phenothiazone, thionol and phenothiazine-5-oxide.

An apparatus and methods are described for the separation of the drug and its metabolites from biological materials and their estimation by colorimetric methods.

The site and form of absorption and oxidation of phenothiazine and the nature and site of the excretory products are discussed.

The chronological distribution of phenothiazine and its metabolites in the chicken and their possible pharmacological significance are considered.

BIBLIOGRAPHY

1. Allen, R. W., Olivier, L., and Peterson, H. O.: The Efficacy of Phenothiazine for the Removal of the Cecal Worm of Chickens. *Vet. Med.* 37 (1942): 412-415.
2. Barnett, E. de B., and Smiles, S.: The Intermolecular Rearrangement of Diphenylamine Ortho-Sulfoxide. *J. Chem. Soc.* 95 (1909): 1253-1266.
3. Benham, G. H.: The Fate of Phenothiazine in Rabbits. I. The Detection of a New Conjugate on Rabbit's Urine After the Feeding of Phenothiazine. *Canad. J. Res.*, 23E, (1945): 71-79.
4. Bernthsen, A.: Zur Kenntniss des Methylenblau and verwanter Farbstoffe. *Berichte der deutschen chemischen Gesellschaft*, 16 (1883): 2896-2911.
5. Bernthsen, A.: Studien in der Methylblaugruppe. *Leibigs Ann. Chemie und Pharm.* 230 (1885): 73-211.
6. Calm, A.: *Berichte der deutschen chemischen Gesellschaft*, 16 (1883): 3799.
7. Campbell, F. M., Sullivan, W. N., Smith, L. E., and Haller, M. L.: Insecticidal Tests of Synthetic Organic Compounds, Chiefly Tests of Sulfur Compounds against Culicine Mosquito Larvae. *Journ. Econ. Ent.*, 27(1934): 1185.
8. Carro, H.: *Engl. Pat.* 3751, from 9.X 1877 and D.R.P. 1886.
9. Clapham, P. A.: On Treating and Preventing Gapeworm Disease. *J. Helmit.*, 24(1950): 53-60.
10. Clare, W. F.: A Photosensitized Keratitis in Young Cattle Following the Use of Phenothiazine as an Anthelmintic. II. The Metabolism of Phenothiazine in Ruminants. *Austral. Vet. J.*, 23, (1947): 340-344.
11. Clare, N. T., Whitten, L. K., and Filmer, D. F.: A Photosensitized Keratitis in Young Cattle Following the Use of Phenothiazine as an Anthelmintic. III. Identification of the Photosensitizing Agent. *Austral. Vet. J.* 23, (1947): 344-348.
12. Collier, H. B., Allen, D. E., and Swales, W. E.: Observation on the Fate of Phenothiazine in Domestic Animals. *Canad. J. Res.*, 21D, (1943): 151-159.

13. De Eds, F., Clyde, W., and Thomas, J. O.: Thionol from Phenothiazine. U. S. 2, 162, 686, June 13.
14. De Eds, F., Eddy, C. W., and Thomas, J. O.: Studies on Phenothiazine in Body. J. Pharmac. Exptl. Therap., 64 (1938): 250-262.
15. De Eds, F., Eddy, C. W., and Thomas, J. O.: Studies on Phenothiazine. V. Fate of Phenothiazine in the Body. J. Pharm., 64, (1938): 250-262.
16. De Eds, F., and Thomas, J. O.: Studies on Phenothiazine. X. The Excretion of Phenothiazine. J. Parasit., 28, (1942): 363-367.
17. De Eds, F., and Thomas, J. O.: Studies on Phenothiazine. IX. The Biliary Excretion and Anthelmintic Action of Thionol. J. Parasit., 27, (1941): 143-151.
18. de Macedo, Luis R. Tavares: "Action of Phenothiazine on Intestinal Protozoa of Turkey and Chickens." Bol. Inst. Vital Brazil, 27 (1944): 47-57.
19. Ellison, T., Todd, A. C., and Harvey, Jr., J.: Isolation and Identification of the Phenothiazine Urinary Conjugate in Dairy Cows. Am. J. Vet. Res., 18, (1957): 956-958.
20. Ellison, T., and Todd, A. C.: The Metabolism of Phenothiazine, N. F., in Dairy Calves. I. Chronological Distribution of Drug in Digestive Tract Contents. Am. J. Vet. Res., 18 (1957): 519-529.
21. Esserman, H. B.: The Mode of Action of Phenothiazine as an Anthelmintic. II. Phenothiazine in the Intestinal Fluid and Nematode Parasites of Treated Animals. Aust. J. Sci. Res., Sec. B., 5 (1952): 485-495.
22. Flatla, J. L.: Phenothiazin (Afitiazin, Fentiazin) som ormeniddel til hest. Norsk. Vet. Tidsskr., 56 (1944): 313-328.
23. Granick, S. and Michaelis, L.: Thionol and its Semiquinone Radical. J. Am. Chem. Soc., 69 (1947): 2983-2986.
24. Griffith, H. J.: A Summary of the Use of Phenothiazine in the Control of Cecal Worms and in the Prevention of Blackhead. Canad. J. Comp. Med., 10 (1946): 218-221.
25. Guilhon, J.: Traitement du Parasitisme Intestinal des Oiseaux par la Thiodiphenylamine. Bull. Acad. Vet. Fr., 18 (1945): 101-111.

26. Guthrie, J. E., and Harwood, P. D.: The Efficacy of Phenothiazine and Nicotine-Bentonite for the Removal of *Heterakis Gallinae* and *Ascaridia Galli* from Chickens. *J. Parasit.*, 28 (Suppl) (1942): 24-25.
27. Harpur, R. P., Denstedt, O. F., and Swales, W. E.: Absorption of Phenothiazine from the Rumen of Sheep, and its Estimation in Blood Serum. *Canad. J. Res.*, D28 (1950): 162-172.
28. Harpur, R. P., Swales, W. E., and Denstedt, O. F.: The Fate of Phenothiazine in the Gastrointestinal Tract of the Sheep. *Canad. J. Res.*, D28 (1950): 143-161.
29. Harwood, P. D.: The Anthelmintic Properties of Phenothiazine. *Exp. Parasit.*, 2(1953): 428-455.
30. Harwood, P. D. and Guthrie, J. E.: The Effect of Nicotine-Bentonite and of Certain Physical States Upon the Efficacy of Phenothiazine Against Nematodes in Fowls. *J. Parasit.*, 30 (1944): 143-152.
31. Harwood, P. D. and Guthrie, J. E.: The Effect of Prickly-ash Bark Upon the Efficacy of Phenothiazine Against Nematodes in Fowls. *J. Parasit.* 30 (1944): 197.
32. Harwood, P. D., Jerstad, A. C., and Swanson, L. E.: The Efficacy of Phenothiazine for the Removal of Ascarids and Nodular Worms from Swine. *J. Parasit.*, 24 (Suppl.) (1938): 16-17.
33. Houston, D. F., Kester, E. B., and De Eds, F.: Phenothiazine Derivatives: Mono-oxygenated Compounds. *J. Am. Chem. Soc.*, 71 (1949): 3816-3818.
34. Houston, D. F., Kester, E. B., and De Eds, F.: Phenothiazine Derivatives: Di-oxygenated Compounds. *J. Am. Chem. Soc.*, 71 (1949): 3819-3822.
35. Knipling, E. F.: Internal Treatment of Animal with Phenothiazine to Prevent Development of Horn Fly Larvae in the Manure. *Journ. Econ. Ent.*, 31 (1938): 315-316.
36. Knoevenagel, E.: Über die Katalytischen Wirkungen des Jods. *Journal für Praktische Chemie*, 89 (1914): 1-50.
37. Krupski, A., und Leemann, W.: Phenothiazin Gegen Darmparasiten der Haustiere. *Schweiz. Arch. Tierheilk.*, 85 (1943): 234-239.
38. Lazarus, M. and Rogers, W. P.: Uptake of Phenothiazine, Labelled with S^{35} , by the Tissues of Nematodes Parasites and their Hosts. *Nature*, 166 (1950): 647-648.

39. Lipson, M.: An Examination of the Urine of Sheep Dosed with Phenothiazine. Austral. J. Exptl. Biol. M. Sci., 18, (1940): 269-272.

40. Lewis, G. N.: J. Am. Chem. Soc., 67 (1945): 770.

41. McCulloch, E. C.: Phenothiazine and Related Products as Controls for Internal Parasites of Poultry. Rep. Wash. Agric. Exp. Sta., 54th, 75, (1943-44).

42. McCulloch, E. C. and Fuller, S. A.: The Efficiencies of Phenothiazine and Related Products in Removing and Controlling Internal Parasites in Poultry. Bull. Wash. St. Agric. Exp. Sta., No. 425, (1942), 81-82.

43. McCulloch, E. C., and Nicholson, L. G.: Phenothiazine for the Removal of Heterakis Gallinae from Chickens. Vet. Med., 35 (1940): 398-400.

44. McCulloch, E. C., and Nicholson, R. C.: The Effect of Phenothiazine on the Hemoglobin Concentration of Chickens. Vet. Med., 37 (1942): 248-252.

45. Meriweather, B.: Phenothiazine vs. Hemoglobin. Vet. Med., 36 (1941): 374-375.

46. Mozolowski, W.: The Quantitative Estimation of Glycuronic Acid and its Conjugated Compounds by Means of the Naphtoresorsinol Test of Tollens. Biochem. J., 34 (1940): 823-828.

47. Nicholson, L. G. and McCulloch, E. C.: Some Effects of Feeding Phenothiazine to Chickens in Various Amounts. J. Am. Vet. Med. Ass., 101 (1942): 205-209.

48. Oliver, W. T.: Studies in Phenothiazine. I. Some in vitro Effects of Phenothiazine on Heterakis Gallinae. Canad. J. Comp. Med., 17 (1953): 435-447.

49. Oliver, W. T., and Funnell, H.: Unpublished data.

50. Olivier, L., Allen, R. W. and Hardcastle, A. B.: Removal of the Cecal Worm, Heterakis Gallinae from Chickens by Feeding Phenothiazine in the Mash. Vet. Med. 38 (1943): 384-386.

51. Oberg, E. and Jones, F. D.: Machinery's Handbook. The Industrial Press, New York, (1954). p. 418.

52. Parish, H. E.: Agric. News Letter (Du Pont), 10 (1942): 35.

53. Pummerer, R. and Gassner, S.: Uber die Desmatropie o.- und p.- Chinoider Salze in der Thiazinreihe. Berichte der deutschen chemischen Gesellschaft, 46 (1913): 2310-2327.

54. Reynoso, R. H.: Parasitosis Intestinal en las aves (Heterakiasis). Alm. Min. Agric., 20 (1945): 331-333.
55. Roberts, F. H. S.: A Preliminary Note on the Efficiency of Phenothiazine Against Some Poultry Helminth. Aust. Vet. J., 16 (1940): 172-174.
56. Schmid, F.: Phenothiazin als Wurmmittle. Dtsch. Tierarztl. Wschr., 50 (1942): 26-28.
57. Schmid, G.: "Treatment of Fowl Coccidiosis with Phenothiazine." Schweiz. Arch. Tierheilk., 88 (1946): 413-417.
58. Sturkie, P. D.: Avian Physiology, Comstock Publishing Assoc., Ithaca, New York (1954): p. 170.
59. Swales, W. E., and Collier, H. B.: Studies on Effects and Excretion of Phenothiazine when Used as an Anthelmintic for Sheep. Canad. J. Res., 18D, (1940): 279-287.
60. Swierstra, D.: "Effect of Phenothiazine on Capillaria Longicollis in Fowl. Tijdschr. Diergeneesk., 73 (1948): 149-153.
61. Temperton, H., and Dudley, F. J.: The Use of Phenothiazine in the Control of Worms Among Laying Pullets. Harper Adams Util. Poult. J., 30 (1945): 37-41.
62. Tobler, J.: Zur Bekämpfung der Geflügelkokzidiose mit Phenothiazin. Schweiz. Arch. Tierheilk., 88 (1946): 458-461.
63. Wasserman, R. H., Trum, B. F., Manroe, R. A., Lane, J. J., and Comar, C. L.: Studies on Depression of Radioiodine by the Thyroid After Phenothiazine Administration. I. Investigations of the Factors in Phenothiazine N. F. Responsible for the Thyroidal Effects. Amer. J. Vet. Res. 17 (1956): 149-152.

Table No. 1

	30 min.*			1 Hour*		
	Ptz.	'zone	Thio.	Ptz.	'zone	Thio.
Crop	835	ND	ND	750	ND	ND
Proventriculus	5.7	ND	ND	7.2	ND	ND
Gizzard	15.2	ND	ND	69.5	ND	ND
Small intestine	ND	ND	ND	3	ND	ND
Caeca	ND	ND	ND	ND	ND	ND
Large intestine	ND	ND	ND	ND	ND	ND
Cloaca	ND	ND	ND	ND	ND	0.2
Excreta	ND	ND	ND	ND	ND	0.3
Bile	ND	ND	ND	ND	ND	ND
Liver	ND	ND**		0.8	0.05**	
Kidneys	ND	ND**		ND	0.1**	
Plasma***	ND	ND**		0.3	0.6**	

Ptz. : Phenothiazine (total milligrams)

'zone: Phenothiazone (total milligrams)

Thio.: Thionol (total milligrams)

ND : Not detected

* Post-treatment

** Sum of total water soluble oxidative derivatives (total milligrams)

*** Plasma: Milligrams per 100 ml.

Table No. 2

	2 Hours*			3 Hours*		
	Ptz.	'zone	Thio.	Ptz.	'zone	Thio.
Crop	584	ND	ND	485	ND	ND
Proventriculus	21.5	ND	ND	1.6	ND	ND
Gizzard	197	ND	ND	250	ND	ND
Small intestine	30	ND	ND	57.2	ND	ND
Caeca	ND	ND	ND	ND	ND	ND
Large intestine	ND	ND	ND	ND	ND	ND
Cloaca	ND	ND	0.24	ND	0.1	0.17
Excreta	ND	ND	0.46	ND	ND	0.6
Bile	ND	ND	ND	ND	ND	ND
Liver	1	0.06**		1.1	0.24**	
Kidneys	ND	0.32**		ND	0.14**	
Plasma***	0.2	1**		0.37	1.1**	

Ptz. : Phenothiazine (total milligrams)

'zone: Phenothiazone (total milligrams)

Thio.: Thionol (total milligrams)

ND : Not detected

* Post-treatment

** Sum of total water soluble oxidative derivatives (total milligrams)

*** Plasma: Milligrams per 100 ml.

Table No. 3

	4 Hours*			5 Hours*			6 Hours*		
	Ptz.	'zone	Thio.	Ptz.	'zone	Thio.	Ptz.	'zone	Thio.
Crop	275	ND	ND	134.5	ND	ND	122.5	ND	ND
Proventriculus	3	ND	ND	1.2	ND	ND	0.8	ND	ND
Gizzard	470	ND	ND	364	ND	ND	197	ND	ND
Small intestine	103	ND	ND	122	ND	ND	330	0.5	ND
Caeca	0.4	0.34		3	0.5	0.24	4.4	1	0.6
Large intestine	0.15	ND	ND	0.2	0.25	0.06	0.2	0.1	0.08
Cloaca	0.5	0.3	0.6	102	2.3	0.6	20	1.6	2.5
Excreta	0.7	0.2	0.9	22	0.6	1.2	115	2.6	4.6
Bile	0.15	ND	ND	0.2	0.5	0.2	0.4	0.65	0.7
Liver	1	0.3**		1.2	0.06**		1	0.06**	
Kidneys	ND	0.22**		ND	0.36**		ND	0.32**	
Plasma***	0.85	2.1**		1	1.9**		2	2.4**	

Ptz. : Phenothiazine (total milligrams)
 'zone: Phenothiazone (total milligrams)
 Thio.: Thionol (total milligrams)
 ND : Not detected

* Post-treatment
 ** Sum of total water soluble oxidative derivatives (total milligrams)
 *** Plasma: milligrams per 100 ml.

Table No. 4

	16 Hours*			24 Hours*			48 Hours*		
	Ptz.	'zone	Thio.	Ptz.	'zone	Thio.	Ptz.	'zone	Thio.
Crop	83	ND	ND	1.3	ND	ND	ND	ND	ND
Proventriculus	0.1	ND	ND	0.6	ND	ND	ND	ND	ND
Gizzard	92	ND	ND	30	ND	ND	6	ND	ND
Small intestine	213	0.85	0.5	112	0.6	0.9	3.5	0.36	0.34
Caeca	5.4	3	0.9	15	1.3	0.64	6.7	1.2	1.02
Large intestine	0.8	0.2	0.15	0.6	0.18	0.13	0.1	1	0.1
Cloaca	50	4.4	1.2	48	1.4	0.25	0.1	1.1	0.1
Excreta	325	13	13.5	552	20	22	642	39	37
Bile	0.12	0.75	0.35	0.25	0.82	0.4	0.06	0.55	0.86
Liver	0.85	0.08**		0.65	0.12**		0.6	0.15**	
Kidneys	ND	0.2**		ND	0.1**		ND	0.16**	
Plasma***	3.5	3.2**		1	2.3**		0.25	0.75**	

Ptz. : Phenothiazine (total milligrams)
 'zone: Phenothiazone (total milligrams)
 Thio.: Thionol (total milligrams)
 ND : Not detected

* Post-treatment
 ** Sum of total water soluble oxidative derivatives
 (total milligrams)
 *** Plasma: milligrams per 100 ml.

Table No. 5

Solvent	Fraction No. 1		Synthetic Phenothiazone	
	Minima at μ	Maxima at μ	Minima at μ	Maxima at μ
Chloroform (Figure No. 8)	500	411-412	500	411-412
Benzene (Figure No. 9)	360;374-378; 485-490	405	360;374-378; 485-490	405
Water (Figure No. 10)	382-386; 522-530	436-437; 660-690	382-384; 522-528	432-438; 660-690
Sodium hydroxide 0.01 N (Figure No. 11)	382-384; 525	435; 670	382-384; 525	435 660-680
d-Tartaric acid, half saturated solution (Figure No. 12)	384-388; 542-548	436-444; 650-654	384-388; 542-546	436-438; 650-655
Acetic acid, 50 per cent solution (Figure No. 13)	380-382; 520	431-434; 650-680	380-382; 522-524	432-436; 650-680

Transmittance Characteristics in Visual Spectral Range of Fraction No. 1 from
Caecal Contents and of Synthetic Phenothiazone, in Various Solvents.

Table No. 6

Solvent	Fraction No. 3		Synthetic thionol	
	Minima at μ	Maxima at μ	Minima at μ	Maxima at μ
Sulfuric acid 8 N (Figure No. 15)	497	705-721	497	690-700
Sodium hydroxide 0.01 N (Figure No. 16)	588	685-700	588	640-700
Water (Figure No. 17)	587	690-700	588	650-690
Ethanol 95P.C. (Figure No. 18)	587	690-695	588	650-690

Transmittance Characteristics in Visual Spectral Range of Fraction No. 3 From Caecal Contents and of Synthetic Thionol, in Various Solvents.

Table No. 7

Solvent	Fraction A		Synthetic phenothiazone	
	Minima at μ	Maxima at μ	Minima at μ	Maxima at μ
Sodium hydroxide 0.01 N (Figure No. 19)	380-382; 520-530	435-440	383-385; 525-530	434-438
Absolute methanol (Figure No. 20)	364-366; 500-505	417-420	364-366; 500-505	417-420

Transmittance Characteristics in Visual Spectral Range of Fraction A from Excreta and of Synthetic Phenothiazone in 0.01 N Sodium Hydroxide and Absolute Methanol.

Table No. 8

Solvent	Minima at μ	
	Fraction B	Synthetic thionol
Absolute methanol (Figures No. 21 and 22)	276-277; 590	277;590
Sodium hydroxide 0.01 N (Figure No. 23)	590	590
Sulfuric acid 8 N (Figure No. 24)	490-497	490-497

Transmittance Characteristics in Ultraviolet and Visual Spectral
Ranges of Fraction B from Excreta and of Synthetic Thionol, in
Various Solvents.

Table No. 9

Solvent	Fraction C		Pure phenothiazine	
	Minima at μ	Maxima at μ	Minima at μ	Maxima at μ
Absolute methanol (Figure No. 25)	253; 316	281-282	253; 316	281-282

Transmittance Characteristics in Ultraviolet Spectral Range
of Fraction C and of Pure Phenothiazine in Absolute Methanol.

Table No. 10

Species	Nature of Urinary Forms	Reference
Rat	Leucophenothiazone and leucothionol	15,16
Rabbit	Leucophenothiazone and glucuronide conjugate of thionol	3,12,15,16,17
Man	Conjugate yielding phenothiazine on acidification	15,16,17
Pig	Conjugate yielding phenothiazine on acidification	10,12
Sheep	Leucophenothiazone, leucothionol, conjugate yielding phenothiazine on acidification and potassium ethereal sulfate of leucophenothiazone	10,12,27,39,59
Calf	Conjugate of phenothiazone and a polypeptide	11
Horse	Leucophenothiazone, leucothionol and ethereal sulfate of leucophenothiazone	12
Dog	Leucophenothiazone and leucothionol	12,17
Cow	Potassium ethereal sulfate of 3-hydroxyphenothiazine	19

Figure No. 1. Chromatographic Chamber

C - Pyrex Cylinder; P_t - Upper Stainless Steel Plate with a Stopcock in its Center; P_b - Lower Stainless Steel Plate; T - Polyethylene Solvent Trough; F - Pyrex Glass Frame; R - Glass Rod.

Figure No. 2. Wooden box for covering chromatographic apparatus.

Figure No. 3. Chromatogram showing the positions, from bottom of phenothiazine, thionol, phenothiazone and phenothiazine-5-oxide respectively.

Figure No. 4 Transmittance curve of the reaction product of phenothiazine-5-oxide; minima at 440 and 520 mμ., maxima at 390 and 470 mμ.

Figure No. 5. Calibration curve of phenothiazine-5-oxide.

Figure No. 6. Ultraviolet absorption of phenothiazine, phenothiazine-5-oxide and phenothiazine-3.

Figure No. 7. Molar extinction coefficient of thionol:

- 2×10^{-5} M and 5×10^{-6} M concentrations in sodium hydroxide 0.01 N,
- 2×10^{-5} M and 5×10^{-6} M concentrations in sulfuric acid 8 N.

Figure No. 8. Spectral curves of Fraction No. 1 extracted from caeca and of synthetic phenothiazone in visual range with chloroform as solvent.

Figure No. 9. Spectral curves of Fraction No. 1 extracted from caeca and of synthetic phenothiazone in visual range with benzene as solvent.

Figure No. 10. Spectral curves of Fraction No. 1 extracted from caeca and of synthetic phenothiazone in visual range with water as solvent.

Figure No. 11. Spectral curves of Fraction No. 1 extracted from caeca and of synthetic phenothiazone in visual range with sodium hydroxide 0.01 N as solvent.

Figure No. 12. Spectral curves in visual range of Fraction No. 1 extracted from caeca and of synthetic phenothiazone with d-tartaric acid, half saturated solution, as solvent.

Figure No. 13. Spectral curves in visual range of Fraction No. 1 extracted from caeca and of synthetic phenothiazone with acetic acid, 50 per cent solution, as solvent.

Figure No. 25. Spectral curves in ultraviolet range of Fraction C extracted from excreta and of the purified phenothiazine with absolute methanol as solvent.

Figure No. 14. Spectral curves in ultraviolet range of Fraction No. 2 extracted from caeca and of purified phenothiazine with absolute methanol as solvent.

Figure No. 15. Spectral curves in visual range of Fraction No. 3
extracted from caeca and of synthetic thionol
with sulfuric acid 8 N as solvent.

Figure No. 16. Spectral curves in visual range of Fraction No. 3
extracted from caeca and of synthetic thionol
with sodium hydroxide 0.01 N as solvent.

Figure No. 18. Spectral curves in visual range of Fraction No. 3
extracted from caeca and of synthetic thionol with
95 per cent ethanol as solvent.

Figure No. 17. Spectral curves in visual range of Fraction No. 3
extracted from caeca and of synthetic thionol with
water as solvent.

Figure No. 19. Spectral curves in visual range of Fraction A
extracted from excreta and of synthetic pheno-
thiazone with sodium hydroxide 0.01 N as solvent.

Figure No. 20. Spectral curves in visual range of Fraction A
extracted from excreta and of synthetic pheno-
thiazone with absolute methanol as solvent.

Figure No. 21. Spectral curves in visual range of Fraction B
extracted from excreta and of synthetic thionol
with absolute methanol as solvent.

Figure No. 22. Spectral curves in ultraviolet range of Fraction B
extracted from excreta and of synthetic thionol
with absolute methanol as solvent.

Figure No. 23. Spectral curves in visual range of Fraction B
extracted from excreta and of the synthetic
thionol with sodium hydroxide 0.01 N as solvent.

Figure No. 24. Spectral curves in visual range of Fraction B
extracted from excreta and of the synthetic
thionol with sulfuric acid 8 N as solvent.

Figure No. 26. (Top) Infra-red transmittance curve of Fraction A,
isolated from excreta. Preparation: mineral
oil paste.

Figure No. 27. (Bottom) Infra-red transmittance curve of synthetic
phenothiazone. Preparation: mineral oil
paste.

Figure No. 28. (Top) Infra-red transmittance curve of Fraction B
isolated from the excreta. Preparation:
mineral oil paste.

Figure No. 29. (Bottom) Infra-red transmittance curve of synthetic
thionol. Preparation: mineral oil paste.

Figure No. 30 Comparison of the spectral curves of purified
phenothiazine and of the fraction extracted
from the liver.

DATE DUE

MAY 19 1960

MAY 25 1960

JUN 9 1960

NOV 29 1960

JUN 14 1962

NOV 30 1962

JUL 4 1963

THESIS:

SF 756.T
P 697
1959

Platonow, Nicolas
Metabolism of
phenothiazine in
chickens.

PLATONOW

SF 756.T
P 697
1959

5595

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
SF 756										TO										P 697										1959										K																																							
FORM										CLASS NO.										DATE (1)										BOOK NO. OR DATE										BOOK NO. OR DATE (2)										COPY																													
FORM										CLASS NO.										DATE (1)										BOOK NO. OR DATE										BOOK NO. OR DATE (2)										COPY																													

THE LIBRARY

UNIVERSITY OF GUELPH

BOOK CARD

- THIS CARD MUST BE KEPT IN THE BOOK POCKET
- THE BORROWER WILL BE RESPONSIBLE IF CARD IS MISSING OR DAMAGED.