

**Title:** The efficacy of teat sealants in dairy cows at dry-off to prevent new intra-mammary infections during the dry-period or clinical mastitis during early lactation: A protocol for a systematic review.

**Authors:** Jan M. Sargeant<sup>1,2</sup>, Charlotte Winder<sup>2</sup>, Annette M. O'Connor<sup>3</sup>, Hannah Wood<sup>4</sup>, Julie M. Glanville<sup>4</sup>, David F. Kelton<sup>2</sup>, Stephen LeBlanc<sup>2</sup>, Todd Duffield<sup>2</sup>.

<sup>1</sup> Centre for Public Health and Zoonoses, University of Guelph, Guelph, ON, Canada, N1G 2W1.

<sup>2</sup> Department of Population Medicine, Ontario Veterinary College, University of Guelph, Guelph, ON, Canada, N1G 2W1.

<sup>3</sup> Department of Veterinary Diagnostic and Production Animal Medicine, Iowa State University College of Veterinary Medicine, Ames, IA

<sup>4</sup> York Health Economics Consortium, University of York, York, YO10 5NQ, United Kingdom

#### **Author contributions:**

All authors contributed to the development of the review question and the methodology described in this proposal. HW and JG developed the search strategy. JMS drafted the protocol, with input and final approval of all co-authors.

#### **Registration:**

This protocol is archived in the University of Guelph's institutional repository (The Atrium; <https://atrium.lib.uoguelph.ca/xmlui/handle/10214/10046>) and published online with Systematic Reviews for Animals and Food (SYREAF) available at: <http://www.syreaf.org/>. The systematic review will be reported using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement guidelines (Liberati et al., 2009). This protocol is reporting using the items (headings) recommended in the PRISMA-P guidelines (Moher et al., 2015).

**Support.** Funding support for this systematic review / meta-analysis / network meta-analysis, including the development of the protocol, was provided by The Pew Charitable Trusts.

#### **Introduction.**

**Rationale:** The majority of antibiotic use in the dairy industry is for intramammary infections (IMI), with a large portion of this aimed at treating and preventing IMI during the dry period (Lam et al., 2102). During dry off, formation of the teat-canal keratin plug plays an important role in susceptibility to IMI (Huxley et al., 2002), but there is wide variation between cows on time to complete closure of the teat-canal, or indeed if closure occurs at all (Dingwell et al., 2003). In heifers, pre-partum IMI is an important risk factor for the development of clinical mastitis in early lactation, and the impact of this disease on future udder health and productivity is far greater than in multiparous animals (Piepers et al., 2009). Moreover, the incidence of clinical mastitis at freshening in heifers is roughly double that of multiparous cows (Ali Naqvi et al., 2018).

Teat sealants provide a non-antibiotic strategy to prevent IMI in the pre-calving period, which is of increasing importance due to concern for antimicrobial use and its relationship with the development of antimicrobial resistance (WHO, 2015). Understanding the efficacy of these products is essential to optimizing their use in order to decrease reliance on antibiotics for both treatment and prevention. Systematic reviews of randomized controlled trials in these areas will yield the highest level of evidence for efficacy of treatment under field conditions (Sargeant and O'Connor, 2014). Establishing the efficacy of teat sealants at dry-off, and pre-partum in heifers, to reduce the incidence of both clinical mastitis and/or IMI, will serve to improve decision makers' ability to engage in effective stewardship of antibiotics through the strategic use of non-antibiotic alternatives.

**Objectives:** The objective of this protocol is to describe the methods for a systematic review – network meta-analyses to address the efficacy of internal or external teat sealants at dry off to prevent new IMI and clinical mastitis early in the subsequent lactation.

The specific PICO elements, which will define the eligibility criteria, are as follows:

- i. *Population:* Pre-partum heifers and dairy cows after their first (or greater) lactation without existing IMI.
- ii. *Intervention:* Internal or external teat sealant given to heifers pre-partum or dairy cows at the time of dry off.
- iii. *Comparator:* No teat sealant, placebo, or an alternative treatment (such as an antibiotic dry-cow preparation).
- iv. *Outcomes:* Critical outcomes will include i) incidence of IMI during the pre-calving period immediately following the intervention, ii) incidence of IMI during the first 30 days of the subsequent lactation and iii) incidence of clinical mastitis during the first 30 days of the subsequent lactation. Secondary outcomes will include total antibiotic use during the first 30 days of the subsequent lactation, milk production during the subsequent lactation, somatic cell count during the first test of the subsequent lactation or the average of the first 3 tests of the subsequent lactation, and the risk of culling due to mastitis during the subsequent lactation.

## Methods

**Eligibility criteria:** In addition to eligibility criteria as described in the PICO elements described above, eligibility criteria will include publication in English. Both published and non-published (grey literature) studies are eligible, provided they report the results of a primary research study with a concurrent comparison group using an eligible study design.

**Study designs eligible:** Controlled trials with natural disease exposure will be eligible. During full-text eligibility screening, we will identify studies that appear to address the review questions but using an observational design or an experimental design with deliberate disease induction; however, these studies will not be included in further steps of the review.

**Information sources:**

We will conduct the literature search in a range of relevant bibliographic databases and other information sources containing both published and unpublished literature. Table 1 presents the resources to be searched.

**Table 1: Databases and information sources to be searched**

Database / information source	Interface / URL
MEDLINE, MEDLINE In-Process and MEDLINE(R) Daily Epub Ahead of Print	Ovid SP
CAB Abstracts	CAB Interface
Science Citation Index	Web of Science
Conference Proceedings Citation Index – Science	Web of Science
Agricola	Proquest

We will also hand-search the table of contents of the following relevant conferences from 1997 to 2018:

- Proceedings of the American Association of Bovine Practitioners;
- World Association for Buiatrics;
- National Mastitis Council Proceedings

The FDA website containing the Freedom of Information New Animal Drug Approvals (NADA) summaries also will be searched.

**Search strategy:**

A Science Citation Index (Web of Science) search strategy designed to identify studies of antibiotic treatments during the dry-off period in dairy cattle is presented in Table 2. The search strategy employs a multi-stranded approach to maximize sensitivity. The conceptual structure is as follows:

- Dairy cattle;
- AND
- Internal or external teat sealants.

**Table 2: Search strategy to identify studies of teat sealants during the dry-off period in dairy cattle in Science Citation Index (Web of Science)**

# 11	#10 AND #4	552
# 10	#9 OR #8 OR #7 OR #6 OR #5	63,874

# 9 TS=(bismuth\* OR Teatseal\* OR Orbeseal\* OR "Orbe-seal\*" OR LockOut\* OR "Lock Out\*" OR Boviseal\* OR "Bovi-seal\*" OR Cepralock\* OR "Cepra-lock\*" OR Noroseal\* OR "Noro-seal\*" OR THexx\* OR "T-Hexx\*" OR Ubroseal\* OR "Ubro-seal\*" OR DryFlex\* OR "Dry-Flex\*" OR StrongHold\* OR "Strong Hold\*") 47,458

# 8 TS=(("teat" OR "teats" OR intramammar\* OR "intra-mammar\*") NEAR/5 barrier\*) 29

# 7 TS=(("teat" OR "teats" OR intramammar\* OR "intra-mammar\*" OR "barrier") NEAR/5 ("dip" OR "dips" OR "dipped" OR "dipping" OR coat\* OR film\*)) 15,018

# 6 TS=((external\* OR internal\* OR persistent\*) NEAR/5 ("seal" OR "seals" OR sealant\* OR "sealed" OR "sealing" OR sealer\* OR plug\*)) 1,007

# 5 TS=(("teat" OR "teats" OR intramammar\* OR "intra-mammar\*" OR "barrier") NEAR/5 ("seal" OR "seals" OR sealant\* OR "sealed" OR "sealing" OR sealer\* OR plug\*)) 590

# 4 #3 OR #2 OR #1 495,342

# 3 TS=(mastiti\* OR ((intramammar\* OR "intra-mammar\*") NEAR/3 (infect\* OR inflamm\*))) 16,589

# 2 TS=(ayrshire\* OR "brown swiss\*" OR "busa" OR "busas" OR canadienne\* OR dexter\* OR "dutch belted\*" OR "estonian red\*" OR fleckvieh\* OR friesland\* OR girolando\* OR guernsey\* OR holstein\* OR illawarra\* OR "irish moiled\*" OR jersey\* OR "meuse rhine issel\*" OR montbeliarde\* OR normande\* OR "norwegian red\*" OR "red poll" OR "red polls" OR shorthorn\* OR "short horn\*") 53,936

# 1 TS=("cow" OR "cows" OR "cattle" OR heifer\* OR "dairy" OR "milking" OR "bovine" OR "bovinae" OR buiatric\*) 465,272

The search strategies will not be limited by date, language, or publication type.

We will conduct searches using each database listed in the protocol, translating the strategy appropriately to reflect the differences in database interfaces and functionality.

### Study records:

**Data management:** We will download the results of searches in a tagged format and load them into bibliographic software (EndNote). The results will be de-duplicated using several algorithms. We will save results from resources that do not allow export in a format compatible with EndNote in Word or Excel documents as appropriate and manually de-duplicate. The de-duplicated search results will be uploaded into online systematic review software (DistillerSR®, Ottawa, ON, Canada). Reviewers will have training in epidemiology and in systematic review methods. Prior to both abstract and full-text screenings, data extraction, and risk of bias assessment, the reviewers assigned to each step will undergo training to ensure consistent data collection using the forms created in DistillerSR®.

**Selection process:** In the first round of screening, abstracts and titles will be screened for eligibility. Two reviewers will independently evaluate each citation for relevance using the following questions:

- 1) Does the study evaluate the use of internal or external teat sealants in pre-partum dairy heifers or at dry-off in dairy cows following the first or greater lactation?  
YES (neutral response), NO (EXCLUDE), UNCLEAR (neutral response)
- 2) Is there a concurrent comparison group? (i.e. controlled trial with natural or deliberate disease exposure or analytical observational study)?  
YES (neutral response), NO (EXCLUDE), UNCLEAR (neutral response)
- 3) Is the full text available in English?  
YES (include for full text screening), NO (EXCLUDE), UNCLEAR (include for full text screening)

Citations will be excluded if both reviewers responded “no” to any of the questions. Any disagreements will be resolved by consensus. If consensus cannot be reached, the article will be marked as “unclear” and will advance to full text screening. A pre-test will be conducted by all reviewers on the first 250 abstracts to ensure clarify of questions and consistency of understanding of the questions.

Following title/abstract screening, eligibility will be assessed through full-text screening. The same questions will be used as for the title / abstract screening. Two reviewers will independently evaluate the full text articles, with any disagreements resolved by consensus. If consensus cannot be reached, a third reviewer will be used.

**Data collection process:** Data will be extracted by two reviewers working independently. Any disagreements will be resolved by consensus or, if consensus cannot be reached, a third reviewer will be used. Authors will not be contacted to request missing data or to clarify published results. A form for data extraction will be created for this review in DistillerSR® and pre-tested on 4 full text articles to ensure question clarity.

#### **Data items:**

Study level data to be extracted include:

- Study design: experimental with natural disease exposure, experimental with deliberate disease exposure (“challenge trial”), or analytical observational
- Country
- Commercial versus research trials
- Year the study was collected
- Months of data collection
- Breed of cattle
- Whether the study population is comprised of pre-partum heifers, first lactation or greater dairy cows or both

- Description of the intervention (specific teat sealant) and, for pre-partum heifers, the time pre-partum when the intervention was applied
- Description of comparison group

The above data will be collect for all of the primary hypothesis-testing studies that are identified as relevant at full text screening (i.e., experimental studies with natural disease exposure, experimental studies with deliberate disease induction, and analytical observational studies). The arm level data, described below, will be extracted only for experimental studies with natural disease exposure.

Arm level data collected:

- Number of animals enrolled
- Number of animals lost to follow up
- Number of animals analyzed
- Any additional concurrent treatments given to the intervention groups – studies with additional treatments will be considered as separate treatments arms to studies with only an internal or external sealant.

#### **Outcomes and prioritization:**

##### ***Critical outcomes (in order of prioritization):***

- Incidence of clinical mastitis during the first 30 days of the subsequent lactation,
- Reduction in new IMI during the dry-cow period or, in pre-partum heifer, the period between treatment and calving,
- Reduction of new IMI during the first 30 days of lactation.

##### ***Secondary outcomes (in order of prioritization):***

- Total antibiotic use to treat clinical mastitis during the first 30 days of the subsequent lactation,
- Milk production during the subsequent lactation,
- Somatic cell count the first test of the subsequent lactation, or the average of the first 3 tests of the subsequent lactation,
- Risk of culling due to mastitis during the subsequent lactation.

These outcomes were prioritized based on their impact on animal health and welfare and their economic importance. Formal evaluation of these criteria for prioritization was not undertaken.

Data will be collected to describe the outcomes that were evaluated for all eligible studies, regardless of study design. The specific outcome data, as described below, will be extracted only for experimental studies with natural disease exposure.

##### ***Outcome data to be collected:***

- 1) Incidence of clinical mastitis during the subsequent lactation
  - a. Case definition of clinical mastitis
  - b. Level at which outcome data were measured (quarter, composite individual)

## 2) Outcomes related to IMI

- a. Method of determining the study subjects were free of IMI at dry-off or, for pre-partum heifer, prior to administering the intervention:
  - i. Negative culture (extract data on quarter or composite)
  - ii. Somatic cell count below a threshold (extract data on threshold and time period for assessment)
  - iii. No clinical case of mastitis during specified duration (extract data on duration)
  - iv. Other (specify)
  - v. Not assessed – excluded from meta-analysis, as cannot distinguish incident from prevalent cases.
- b. Level at which the outcomes were measured (quarter, composite individual, group)
- c. Method of diagnosis of IMI status
  - i. Number of milk samples used to classify IMI status and timing of sample collection if > 1
  - ii. Whether National Mastitis Council (NMC) Laboratory Methods were stated as used
  - iii. If other methods were used in parallel or exclusively e.g. PCR; Petrifilm or selective media
- d. Type of bacteria:
  - i. Individual bacteria results will be extracted for: Coliforms, Strep. uberis, Strep. agalactica, Staph. aureus
  - ii. Grouped bacteria results will be extracted for: Major contagious mastitis pathogens (Staph. aureus and Strep. agalactia), and Major environmental mastitis pathogens (Strep. uberis and coliforms)

For each of the primary and secondary outcomes, we will extract the possible metrics in the following order:

- 1<sup>st</sup> priority: Adjusted summary effect size (<sub>adjusted</sub> risk ratio or <sub>adjusted</sub> odds ratio, mean differences for continuous outcomes) and variables included in adjustment and corresponding precision estimate
- 2<sup>nd</sup> priority: Unadjusted summary effect size
- 3<sup>rd</sup> priority: Arm level risk of the outcome, or arm level mean of the outcome (continuous outcomes)
- Variance components.

If variance estimates are not reported, but the authors provide the data necessary to calculate them using standard formulas, we will calculate these data. If results are provided only in graphical form, we will estimate the numerical results using WebPlotDigitizer

(<https://automeris.io/WebPlotDigitizer/>), if the full text is in a suitable format for using this resource.

**Risk of bias in individual studies:** Risk of bias will only be assessed for controlled trials with natural disease exposure. Risk of bias assessment will be performed at the outcome level for each of the critical outcomes using the Cochrane risk of bias instrument (Higgins et al., 2016), with the signaling questions modified as necessary for the specific review question. The ROB-2.0 for RCTs will be used. This tool is available at <https://sites.google.com/site/riskofbiastool/welcome/rob-2-0-tool>.

### **Data synthesis:**

**Network meta-analysis.** Network meta-analysis (aka mixed treatment comparison meta-analysis) will be conducted for each of the primary outcomes. Network meta-analysis will use the approach described by NICE Decision Support Unit technical document (Dias et al., 2014; O'Connor et al., 2013, O'Connor et al., 2016). The approach to reporting will use the PRISMA-NMA (<http://www.prisma-statement.org/Extensions/NetworkMetaAnalysis.aspx>). Planned a priori sub-group analyses will be conducted for randomized versus non-randomized trials, heifers versus first lactation or older cows, and internal versus external sealants.

**Meta-bias(es):** Small study effects (“publication bias”) will be assessed for all antibiotic-comparator combinations where there are at least 10 studies in the meta-analysis. If feasible, we will use approaches to assessing publication bias in the network of evidence using previously proposed approaches (Mavridis et al., 2013; Mavridis et al., 2014).

**Confidence in cumulative evidence:** The quality of evidence for each critical outcome will be assessed using the approach proposed by GRADE (GRADE, 2015, Puhan et al., 2014), while also considering the nature of the network meta-analysis (Jansen et al., 2011)

### **Discussion:**

This systematic review will provide a synthesis of the current evidence regarding the efficacy of teat sealants used pre-partum in heifers and at dry off in dairy cows to prevent IMI and clinical mastitis. Results will be helpful for veterinarians and dairy producers when making evidence-informed decisions regarding dry cow management to reduce mastitis and potentially reduce the need to use antibiotics in dairy cows at dry off or to treat clinical disease. The results also will be helpful for identifying specific gaps in knowledge related to the efficacy of these products for targeting additional research.



## References:

- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. 2009. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol.* 62: e1-e34. [10.1016/j.jclinepi.2009.06.006](https://doi.org/10.1016/j.jclinepi.2009.06.006).
- Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. 2015. Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) 2015 statement. *Syst Rev.* 4(1):1. [doi: 10.1186/2046-4053-4-1](https://doi.org/10.1186/2046-4053-4-1)
- Lam, T.J.G.M., E. van Engelen, C.G.M. Scherpenzeel, J.J Hage. 2012. Strategies to reduce antibiotic usage in dairy cattle in the Netherlands. *Cattle Practice* 20:166-171.
- Huxley, J.N., M.J. Green, L.E. Green, A.J. Bradley. 2002. Evaluation of the efficacy of an internal teat sealer during the dry period. *J. Dairy Sci.* 85: 551-561.
- Dingwell, R.T., D.F. Kelton, K.E. Leslie. 2003. Management of the dry cow in control of peripartum disease and mastitis. *Vet. Clin. Food Anim.* 19: 235-265.
- Piepers, S., S. De Vliegher, A. De Kruif, G. Oposmer, H.W. Barkema. 2009. Impact of intramammary infections in dairy heifers on future udder health, milk production, and culling. *Vet. Microbiol.* 134:113-120.
- Ali Naqvi, S., J. De Buck, S. Dufour, H.W. Barkema. 2018. Udder health in Canadian dairy heifers during early lactation. *J. Dairy Sci.* 101:3233-3247.
- World Health Organization. 2015. Global action plan on antimicrobial resistance. Accessed June 7, 2018. [http://www.who.int/iris/bitstream/10665/193736/1/9789241509763\\_eng.pdf?ua=1](http://www.who.int/iris/bitstream/10665/193736/1/9789241509763_eng.pdf?ua=1)
- Sargeant, J.M., and A.M. O'Connor. 2014. Introduction to systematic reviews in animal agriculture and veterinary medicine. *Zoon. Public Health.* 61: 3 – 9.
- Higgins J, Sterne J, Savović J, Page M, Hróbjartsson A, et al., 2016. A revised tool for assessing risk of bias in randomized trials In: Chandler J, McKenzie J, Boutron I, Welch V (editors). *Cochrane Methods. Cochrane Database of Systematic Reviews Issue 10 (Suppl 1)*. [dx.doi.org/10.1002/14651858.CD201601](https://dx.doi.org/10.1002/14651858.CD201601).
- Dias, S., N.J. Welton, A.J. Sutton, A.E. Ades. 2014. NICE DSU technical support document 2: A generalized linear modeling framework for pairwise and network meta-analysis of randomized controlled trials. Decision Support Unit. Accessed Dec. 1 2017. <https://www.ncbi.nlm.nih.gov/pubmedhealth/n/nicedsutsd2/pdf/>

O'Connor AM, Coetzee JF, da Silva N, Wang C. 2013. A mixed treatment comparison meta-analysis of antibiotic treatments for bovine respiratory disease. *Prev Vet Med* 110:77-87.

O'Connor AM, Yuan C, Cullen JN, Coetzee JF, da Silva N, Wang C. 2016. A mixed treatment meta-analysis of antibiotic treatment options for bovine respiratory disease - An update. *Prev Vet Med.* 132:130-9.(doi):10.1016/j.prevetmed.2016.07.003.

Mavridis D, Sutton A, Cipriani A, Salanti G. 2013. A fully Bayesian application of the Copas selection model for publication bias extended to network meta-analysis. *Stat Med* 32:51-66.

Mavridis D, Welton NJ, Sutton A, Salanti G. 2014. A selection model for accounting for publication bias in a full network meta-analysis. *Stat Med* 33:5399-5412.

GRADE. 2015. A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis. *BMJ* 350:h3326.

Puhan MA, Schunemann HJ, Murad MH, Li T, Brignardello-Petersen R, Singh JA, Kessels AG, Guyatt GH, Group GW. 2014. A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis. *BMJ* 349:g5630.

Jansen JP, Fleurence R, Devine B, Itzler R, Barrett A, Hawkins N, Lee K, Boersma C, Annemans L, Cappelleri JC. 2011. Interpreting indirect treatment comparisons and network meta-analysis for health-care decision making: report of the ISPOR Task Force on Indirect Treatment Comparisons Good Research Practices: part 1. *Value in health : the Journal of the International Society for Pharmacoeconomics and Outcomes Research* 14:417-428.